Western Water Assessment | CIRES | University of Colorado Boulder

http://wwa.colorado.edu

Jeff Deems
Western Water Assessment
National Snow and Ice Data Center
University of Colorado

Tom Painter
NASA Jet Propulsion Laboratory

and the ASO Team

The NASA JPL Airborne Snow Observatory

Snow Water Equivalent

Riegl Q1560 ALS
800 kHz Pulse Rate
2 lasers with fore/aft pointing

Snow Albedo

CASI-1500 Imaging Spectrometer
0.35-1.05 µm
2m spatial resolution from 4000m

aso.jpl.nasa.gov

- Retrieve topography: snow-free & snow-on
- Difference gives snow depth
- SWE from assimilation of modeled density field constrained by observations
- SWE variation primarily due to depth

Snow depth & SWE from LiDAR

- majority of spatial variability in SWE is due to snow depth
- depth can be measured by differential elevation mapping
  - collect snow-free & snow-covered data sets
  - classify & remove vegetation
  - subtract snow-free from snow-covered
- apply obs/modeled density
  - SWE = depth * density
  - SNOTEL/manual obs + snow model can estimate density well



LiDAR-derived snow depth, Colorado (Deems et al., 2013)



SNOWPACK model, courtesy CAIC

Snow Albedo from Hyperspectral Imagery



Riegl Q1560 ALS

- Twin lasers
- Long range
  - 5000m+ AGL
- Fore/aft pointing enhances coverage & geometry in steep terrain





- Wide FOV allows large overlap or efficient planning
- Full waveform processing improves subcanopy surface detection

ASO Compute System

operational products

- 24-hour data product latency
- products:
  - snow depth
  - SWE (using snow pillow + modeled densities)
  - broadband albedo
- LiDAR/spectrometer workflows are interdependent
  - LiDAR DSM used for terrain correction
  - spectrally snow-free areas forced to zero snow depth



ASO Demonstration Missions 1 – 3 2013/14/15

2 Primary study basins:

- Tuolumne River, Yosemite National Park, CA
- Uncompahgre River, CO





Snow-off August, 2012

Snow-on Spring 2013-2015

- weekly flights in CA
- monthly flights in CO

Example Results: 2014 & 2015

Tuolumne Basin, CA – Yosemite National Park

Tuolumne River Basin SWE - 2014



Tuolumne River Basin SWE - 2014



Tuolumne River Basin SWE - 2014



Tuolumne River Basin SWE - 2014



Tuolumne River Basin SWE - 2014



Tuolumne River Basin SWE - 2014



Tuolumne River Basin SWE - 2014



Tuolumne River Basin SWE - 2014



Tuolumne River Basin SWE - 2014



Tuolumne River Basin SWE - 2014



Tuolumne River Basin SWE - 2014



Tuolumne River Basin SWE - 2015

Tuolumne River Basin SWE - 2015



Tuolumne River Basin SWE - 2015



Tuolumne River Basin SWE - 2015



Tuolumne River Basin SWE - 2015



Tuolumne River Basin SWE - 2015



Tuolumne River Basin SWE - 2015



Tuolumne Basin SWE & Reservoir Inflow



Banded SWE Report

Uncompahgre River above Ridgway, 30 April 2015



Applications

Integrating ASO with water management…

Hetch Hetchy Reservoir Operations

balancing streamflow, water supply, & hydropower needs











Tuolumne Basin Hydrology Modeling

SWE/Met Stations & PRMS Model Units



ASO SWE & Albedo aggregated to PRMS hydrologic response units



Tuolumne Basin Hydrology Modeling

Results: 2014



Courtesy Bruce McGurk – McGurk Hydrologic

Current ASO products

LiDAR

- High resolution DEM & DSM
- Tree/canopy height maps
- Terrain morphology
- Water body mask
- Surface roughness
- Snow depth
- SWE
- SCA and SC%
- Elevation band SWE map & report
- SWE time series
- Snowline estimation

Spectrometer

- Land surface albedo
- High resolution imagery (RGB)
- High resolution land cover classification
- Snow albedo





SWE & SCA Reports

- Integration with existing data needs
- Banded SWE products
  - Map & tabular
- Basin aggregate
  - SWE & SCA



KC20150531_bandedSWE.txt

```
Airborne Snow Observatory, Jet Propulsion Laboratory, NASA
Kings Basin - full basin data (North & South)
Processing Date: Fri Jun 05 19:55:28 2015
Flight Code and Date: KC20150531
Exported by swe_at_9_elev_bands.pro
SWE file=U:\ASO_Flight_Data\snowon_2015\LIDAR\processed\USCAKC20150531f1a1\KC20150531_SUPERswe_50p0m_agg
Band 1 =       3000.00 to       4000.00 ft
Band 2 =       4000.00 to       5000.00 ft
Band 3 =       5000.00 to       6000.00 ft
Band 4 =       6000.00 to       7000.00 ft
Band 5 =       7000.00 to       8000.00 ft
Band 6 =       8000.00 to       9000.00 ft
Band 7 =       9000.00 to       10000.0 ft
Band 8 =       10000.0 to       11000.0 ft
Band 9 =       11000.0 to       60000.0 ft
```

Band,	total area (sq.mi),	snow area (sq.mi),	band coverage (%),	SWE volume (AF),	Mean SWE depth (inches)
Band1	0.0463	0.0000	0.0	0.0000	0.0000
Band2	7.9016	0.0000	0.0	0.0000	0.0000
Band3	26.5879	0.0512	0.2	2.6818	0.9829
Band4	48.4558	0.1959	0.4	4.7924	0.4586
Band5	73.4540	0.3745	0.5	5.4774	0.2742
Band6	106.8306	0.5039	0.5	7.3039	0.2718
Band7	130.6878	3.1535	2.4	111.7572	0.6645
Band8	176.7517	34.4596	19.5	3695.4087	2.0107
Band9	251.1255	103.3112	41.1	19519.0371	3.5425

TB20150528_swe.report_OUT.txt

```
calculate basin water volume in snowpack
*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*
Tuolumne River Basin
Airborne Snow Observatory
Date: 05/28/2015
surveyed area:                                  1175.54
snow covered area (km^2):                       396.557
percent snow coverage :                         33.7342
volume of snowpack water (m^3):            5.34373e+007
volume of snowpack water (acre-ft):             43322.3
--  --  --  --  --  --  --  --  --  --  --  --  --  --
mean swe (m):                                  0.134753
mean depth (m):                                0.283495
max depth (m):                                  4.60309
min rho  (kg/m^3):                              218.189
max rho  (kg/m^3):                              591.896
mean rho (kg/m^3):                              485.464
*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*^*
file used:
U:\ASO_Flight_Data\snowon_2015\LIDAR\processed
\USCATB20150528f1a1\TB20150528_SUPERswe_50p0m_agg
```

ASO Expansion: California

State of CA & water manager support
for full Sierra Nevada coverage





ASO Expansion: Colorado River Basin

- Uncompahgre River
- Grand Mesa
- Rio Grande/Conejos R.
  (Colorado Water Conservation Board)
- Wyoming:
  (State Engineer's Office)
  - Wyoming Range
  - Wind River Range
  - Sierra Madre
  - Medicine Bow Mountains



Megafire Disaster Response: NASA JPL and USDA Forest Service inter-agency collaborations

- Objective: characterize pre, active and post-burn conditions of megafires, to observe ecosystem properties influencing fire probability, behavior and recovery as a basis for aiding management
- Coverage: California King (2014) and Rim (2013) megafires
- Data collection, processing and dissemination (wildfire.jpl.nasa.gov) is a collaborative effort that aids disaster response and post-fire recovery planning such as:
  - Identify endangered species habitat
  - Water quality assessment
  - Erosion assessment
  - Removal of hazardous logs
  - Timber harvesting



LiDAR before (left) and after (right) King Fire

AVIRIS hyperspectral improves burn area imaging over Rim Fire



A) MASTER – 1 thermal infrared band, a proxy for NIROPS (~ 1 m resolution) used in active fire management perimeter mapping

B) NIROPS progression with MODIS multi-band thermal infrared active fire pixels (~ 1000 m resolution) can be used to calculate Fire Intensity

C) MASTER multiband thermal infrared (35 m resolution)



New Opportunities in Water Management…

snow & weather patterns are increasingly different from past years

- dust on snow
- climate warming

new tools are becoming available to support resilient management of water resources

- expanded measurements of the snow energy balance
- *airborne & satellite mapping of snow depth & albedo*
- high resolution weather forecasts
- integrated snowmelt & hydrology forecasts

Contact:

Jeff Deems
deems@nsidc.org

Tom Painter
thomas.painter@jpl.nasa.gov